## Complete Summary

#### TITLE

Venous thromboembolism (VTE): percentage of adult patients suspected of deep vein thrombosis (DVT) who have leg duplex ultrasound with compression performed despite a low clinical pretest probability and a negative D-dimer test.

## SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Venous thromboembolism. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2006 Feb. 91 p. [202 references]

#### Measure Domain

#### PRIMARY MEASURE DOMAIN

#### **Process**

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the Measure Validity page.

### SECONDARY MEASURE DOMAIN

Does not apply to this measure

#### **Brief Abstract**

## **DESCRIPTION**

This measure is used to assess the percentage of adult patients suspected of deep vein thrombosis (DVT) who have leg duplex ultrasound with compression performed despite a low clinical pretest probability and a negative D-dimer test.

#### **RATIONALE**

The priority aim addressed by this measure is to improve quality of care and costeffectiveness of the diagnosis and treatment of venous thromboembolism (VTE).

#### PRIMARY CLINICAL COMPONENT

Venous thromboembolism (VTE); deep vein thrombosis (DVT); duplex ultrasound with compression; low clinical pretest probability; D-dimer test

#### DENOMINATOR DESCRIPTION

Total number of adult patients suspected of deep vein thrombosis (DVT) who have a low clinical pretest probability and a negative D-dimer test (see the related "Denominator Inclusions/Exclusions" field in the Complete Summary)

#### NUMERATOR DESCRIPTION

Total number of adult patients suspected of deep vein thrombosis (DVT) who have a low clinical pretest probability and a negative D-dimer test who undergo a leg duplex ultrasound with compression

#### Evidence Supporting the Measure

#### EVIDENCE SUPPORTING THE CRITERION OF QUALITY

 A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

#### NATIONAL GUIDELINE CLEARINGHOUSE LINK

Venous thromboembolism.

#### Evidence Supporting Need for the Measure

## NEED FOR THE MEASURE

Unspecified

#### State of Use of the Measure

STATE OF USE

Current routine use

**CURRENT USE** 

Internal quality improvement

#### Application of Measure in its Current Use

CARE SETTING

Physician Group Practices/Clinics

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians

# LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED **Group Clinical Practices** TARGET POPULATION AGE Age greater than or equal to 18 years TARGET POPULATION GENDER Either male or female STRATIFICATION BY VULNERABLE POPULATIONS Unspecified INCIDENCE/PREVALENCE Unspecified ASSOCIATION WITH VULNERABLE POPULATIONS Unspecified **BURDEN OF ILLNESS** Unspecified **UTILIZATION** Unspecified COSTS Unspecified IOM CARE NEED Getting Better IOM DOMAIN

Effectiveness

#### Data Collection for the Measure

#### CASE FINDING

Users of care only

#### DESCRIPTION OF CASE FINDING

Adults 18 years and older evaluated for possible deep vein thrombosis (DVT)

A list of all adult patients evaluated for DVT during the previous target period. The medical records can be reviewed to determine the documentation of a low clinical pretest probability, negative D-dimer test and the performance of a leg duplex ultrasound with compression

Data may be collected semiannually.

#### DENOMINATOR SAMPLING FRAME

Patients associated with provider

#### DENOMINATOR INCLUSIONS/EXCLUSIONS

#### Inclusions

Total number of adult patients suspected of deep vein thrombosis (DVT) who have a low clinical pretest probability\* and a negative D-dimer test

## \*Wells Model of Clinical Pretest Probability of DVT

- Active Cancer (current treatment or treated within the last 6 months, or palliative) -- 1 point
- Paralysis, paresis, or recent plaster immobilization of lower extremity -- 1 point
- Recently bedridden for more then 3 days or major surgery within 4 weeks -- 1 point
- Localized tenderness along the distribution of the deep venous system -- 1 point
- Entire leg swollen -- 1 point
- Calf swollen by more then 3 cm when compared to asymptomatic leg (measured 10 cm below the tibial tuberosity) -- 1 point
- Pitting edema (greater in the symptomatic leg) -- 1 point
- Collateral superficial veins (non-varicose) -- 1 point
- Alternative diagnosis as likely or greater than that of deep vein thrombosis 2 points

## Scoring

If both legs are symptomatic, score the more severe side

High risk scored 3 or more

- Moderate risk scored 1-2
- Low risk scored 0 or less

Exclusions Unspecified

#### RELATIONSHIP OF DENOMINATOR TO NUMERATOR

All cases in the denominator are equally eligible to appear in the numerator

## DENOMINATOR (INDEX) EVENT

Clinical Condition
Diagnostic Evaluation

#### DENOMINATOR TIME WINDOW

Time window precedes index event

#### NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

Total number of adult patients suspected of deep vein thrombosis (DVT) who have a low clinical pretest probability and a negative D-dimer test who undergo a leg duplex ultrasound with compression

Exclusions Unspecified

## MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

## NUMERATOR TIME WINDOW

Episode of care

DATA SOURCE

Medical record

#### LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

#### Computation of the Measure

**SCORING** 

Rate

INTERPRETATION OF SCORE

Better quality is associated with a lower score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

Internal time comparison

#### **Evaluation of Measure Properties**

#### EXTENT OF MEASURE TESTING

Unspecified

#### Identifying Information

#### ORIGINAL TITLE

Percentage of adult patients suspected of DVT who have leg duplex ultrasound with compression performed despite a low clinical pretest probability and a negative D-dimer test.

MEASURE COLLECTION

Venous Thromboembolism Measures

DEVELOPER

Institute for Clinical Systems Improvement

**ADAPTATION** 

Measure was not adapted from another source.

RELEASE DATE

#### 2006 Feb

#### **MEASURE STATUS**

This is the current release of the measure.

#### SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Venous thromboembolism. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2006 Feb. 91 p. [202 references]

#### MEASURE AVAILABILITY

The individual measure, "Percentage of adult patients suspected of DVT who have leg duplex ultrasound with compression performed despite a low clinical pretest probability and a negative D-dimer test," is published in "Health Care Guideline: Venous Thromboembolism." This document is available from the <a href="Institute for Clinical Systems Improvement (ICSI)">Institute for Clinical Systems Improvement (ICSI)</a> Web site.

For more information, contact ICSI at, 8009 34th Avenue South, Suite 1200, Bloomington, MN 55425; phone: 952-814-7060; fax: 952-858-9675; Web site: www.icsi.org; e-mail: icsi.info@icsi.org.

#### NQMC STATUS

This NQMC summary was completed by ECRI on May 1, 2006.

#### COPYRIGHT STATEMENT

This NQMC summary (abstracted Institute for Clinical Systems Improvement [ICSI] Measure) is based on the original measure, which is subject to the measure developer's copyright restrictions.

The abstracted ICSI Measures contained in this Web site may be downloaded by any individual or organization. If the abstracted ICSI Measures are downloaded by an individual, the individual may not distribute copies to third parties.

If the abstracted ICSI Measures are downloaded by an organization, copies may be distributed to the organization's employees but may not be distributed outside of the organization without the prior written consent of the Institute for Clinical Systems Improvement, Inc.

All other copyright rights in the abstracted ICSI Measures are reserved by the Institute for Clinical Systems Improvement, Inc. The Institute for Clinical Systems Improvement, Inc. assumes no liability for any adaptations or revisions or modifications made to the abstracts of the ICSI Measures.

## © 2006 National Quality Measures Clearinghouse

Date Modified: 9/25/2006

## FIRSTGOV

